Effects of cholinesterase inhibitors

If acetylcholine was released at locations where it acted on sympathetic nerve terminals, then drugs that affected cholinergic mechanisms should have secondary effects arising from the way in which actions of the acetylcholine on sympathetic noradrenergic transmission were affected. A number of observations designed to investigate the cholinergic link hypothesis of Burn & Rand (see Introduction for references) has been carried out. Only those in which anticholinesterase drugs were used will be reviewed briefly The expectation of the cholinergic link hypothesis is that anticholinesterases might enhance noradrenergic transmission. the light of other knowledge, there is the possibility that there would be inhibition of noradrenergic transmission mediated by acetylcholine acting on prejunctional cholinoceptors. In most of the experiments concerned with the cholinergic link hypothesis, atropine was used to block the postjunctional action of acetylcholine. However, potentiation of the action of acetylcholine by inhibiting enzymatic hydrolysis might overcome blockade of muscarinic cholinoceptors thereby confounding the findings of such experiments.

Inhibitors of cholinesterase enhanced responses to sympathetic nerve stimulation in rabbit uterus (Varagic, 1956) and isolated heart (Hukovic, 1966), guinea-pig vas deferens (Boyd, Chang & Rand, 1960; Burn & Weetman, 1963) and taenia coli (Ng, 1966), dog femoral artery in situ (Bernard & de Schaepdryver, 1964) and retractor penis (Armitage & Burn, 1967), perfused rabbit ear (Burn & Rand, 1960a) and rabbit ear artery (Rand & Varma, 1970) and rat mesenteric vessels (Malik, 1970). However, the enhancing effect reported on the cat nictitating membrane by Bacq & Frederiq (1935) and by Burn, Rand & Wein (1963) could not be repeated by Mirkin & Cervoni (1962), Gardiner, Hellmann & Thompson (1962) or Bowman, Callingham & Cuthbert (1964), but see Burn & Rand (1965). Furthermore, no enhancing effect could be detected in cat spleen (Blakeley et al., 1963; Thoenen, et al., 1966), rabbit ear artery (Hume et al., 1972) and rabbit pulmonary artery (Bevan & Su, 1964). To complicate matters further, responses to sympathetic nerve stimulation were reduced by cholinesterase inhibitors in cat pilomotor muscles (Hellmann, 1963), dog gastric artery (Van Hee & Vanhoutte, 1978) and retractor penis muscles of various species (Klinge & Sjöstrand, 1977).

Effect of acetylcholine released from parasympathetic nerves on noradrenergic transmission

The tissues with a double supply comprising cholinergic parasympathetic and noradrenergic sympathetic inputs include the heart, the iris, the salivary glands, the gastrointestinal tract and pelvic viscera. Of these, the heart is perhaps the most easily studied: in any event, it has been the most studied.

Acetylcholine released by vagus nerve stimulation in the isolated perfused rabbit heart inhibited the release of noradrenaline in response to sympathetic nerve stimulation in the rabbit isolated heart (Löffelholz & Muscholl, 1970; Langley & Gardiner, 1977; Muscholl et al., 1979), but not in the isolated perfused chicken heart (Löffelholz, 1975).

In field stimulated atria, both acetylcholine and noradrenaline would be released, and in rabbit atria, the released acetylcholine inhibited noradrenaline release since atropine increased, and physostigmine decreased the stimulation-induced release of noradrenaline and the positive inotropic response (Hope et al., 1974; Muscholl & Muth, 1982). Similarly, in dog papillary muscle, the positive inotropic response to sympathetic nerve stimulation was enhanced by atropine (Endoh & Hashimoto, 1970). However, these effects were not demonstrable in guinea-pig atria, on which exogenous acetylcholine had no effect (Hope et al., 1974; Story et al., 1975), or in rat atria, in which exogenous acetylcholine did inhibit noradrenaline release (Loiacono & Story, unpublished observations).

Observations on the dog heart in situ have demonstrated the importance of cholinergic modulation of cardiac sympathetic tone by inhibition of noradrenaline release in addition to the direct actions of acetylcholine on the pacemaker cells and the contractile activity of cardiac myocytes. Levy & Blattberg (1976) and Lavallée et al. (1978) showed that stimulation of the vagus nerves decreased the release of noradrenaline as well as the positive inotropic response to sympathetic nerve stimulation in the dog heart.

The complex interaction between the cholinergic and noradrenergic mechanisms in the control of cardiac activity was first noted by Rosenblueth & Simeone (1934), when they observed that the extent of the bradycardia produced by vagal stimulation was greater in the presence than in the absence of sympathetically maintained tachycardia. Similar observations were made by other workers (Samaan, 1935; Levy & Zieske, 1969; Warner & Russell, 1969). If the parameters of stimulation of the vagus and sympathetic nerves to the dog heart are adjusted so that the responses to each separately are equal in magnitude but opposite in direction, then when the two nerves are stimulated simultaneously the heart rate decreases as if there were no sympathetic stimulation (Levy & Zieske, 1969). This effect has been termed 'accentuated antagonism' (Levy, 1971; Levy & Martin, 1979; Stuesse, Wallich & Levy, 1979), signifying that the resultant effect is not merely the algebraic sum of the two separate responses of bradycardia and tachycardia to vagal and accelerans nerve stimulation, respectively; in fact, the bradycardia predominates (Levy, 1984). This phenomenon also occurs when the pacemaker is in the atrioventricular bundle rather than in the sinoatrial node (Wallick, Felder & Levy, 1978). In contrast, the effects of vagal and sympathetic nerve stimulation on atrioventricular conduction time are simply additive (Levy & Ziescke, 1969; Wallick, Felder & Levy, 1982).

The increase in heart rate produced in the conscious dog by methylatropine (a quaternary analogue of atropine that does not enter into the brain) is due to blockade of the inhibitory effect of acetylcholine released by tonic activity of the vagus nerves on prejunctionmal muscarinic cholinoceptors of noradrenergic terminals, resulting in a greater release of noradrenaline from the tonically active sympathetic nerves (Lokhandwala, Cavero & Buckley, 1973).

The effects on myocardial contractility of cholinergic and noradrenergic mechanisms, like those on chronotropic activity,

display a complex interaction. The effect of acetylcholine on ventricular contractility in the dog is a small reduction; however, when the sympathetic nerves are stimulated or a noradrenaline infusion is given, the negative inotropic effect of acetylcholine is much greater (Hollenberg, Carriere & Barger, 1965). Similar observations on the negative inotropic responses to vagal stimulation were made by Levy, Ng, Martin & Zieske (1966): in their studies, vagal stimulation alone was virtually without effect on ventricular contractility; however, when the force of contraction was increased by sympathetic nerve stimulation, vagal stimulation decreased the force almost to the prestimulation level. The effects of vagal and sympathetic stimulation on the force of atrial contractions exhibited similar interactions (Stuesse et al., 1979).

The evidence indicates clearly that the interaction between the cholinergic and noradrenergic innervations of the heart are not confined to their opposing postjunctional actions but extend to prejunctional interactions (Levy, 1971, 1984; Higgins, Vatner & Braunwald, 1973; Story et al., 1975). The postjunctional interactions have been described by Watanabe, Lindemann & Fleming, (1984).

In addition to cholinergically induced inhibition of noradrenergic transmission, the reverse interaction, noradrenergically induced inhibition of cholinergic transmission has been reported (Vizi & Knoll, 1971; Story et al., 1975; see also: Wetzel, Goldstein & Brown, 1985; Loiacono & Story, 1986). Neuronal interactions may also involve peptide cotransmitters in addition to the classical transmitters of autonomic nerves. Thus, Potter (1985, 1987) showed that stimulation of cardiac sympathetic nerves in the dog resulted in a prolonged inhibition of the response to vagus nerve stimulation that was not due to noradrenaline but was mimicked by neuropeptide Y. It remains to be shown whether there is a peptide cotransmitter in cardiac cholinergic nerves and, if so, whether it affects noradrenergic transmission.

Effects of acetylcholine released from sympathetic cholinergic nerves on noradrenergic transmission

The two tissues to be considered in this section are the vas deferens and blood vessels. They are generally thought to be innervated by sympathetic nerves, except for some blood vessels in certain specialised regions such as the blush areas and erectile tissue which also receive a parasympathetic innervation. Nevertheless, they have a cholinergic as well as a noradrenergic innervation. In the case of the vas deferens it is not clear whether there are separate cholinergic nerves or whether the noradrenergic nerves contain a cholinergic component.

Vas deferens

Prejunctional muscarinic cholinoceptors may be involved in the neuronal control of the activity of the guinea-pig vas deferens, in which muscarinic agonists inhibit stimulation-induced noradrenaline release (Stjärne, 1975; Leighton & Westfall, 1976). There appear to be cholinergic nerves in this tissue since it contains acetyltransferase (Ohlin & Strömblad, 1963) and field stimulation releases acetylcholine (Knoll, Somogyi, Illes & Vizi, 1972).

If endogenously released acetylcholine had an inhibitory effect on stimulation-induced noradrenaline release, blockade of the prejunctional muscarinic cholinoceptors should enhance noradrenaline release. This does not appear to be the case in most tissues (see Starke, 1977). However, there are exceptions, as mentioned above for rabbit atria, and here for the vas deferens.

The endogenously released acetylcholine apparently inhibits noradrenaline release since the amount of noradrenaline released by field stimulation is increased by atropine (Stjärne, 1975). Paradoxically, muscarinic agonists enhance the contractile response to field stimulation (Sjöstrand, 1961; Sjöstrand & Swedin, 1968; Lamble & Lamble, 1975), but this is presumably by a postjunctional action: whether it involves interaction with the contractile effect of noradrenaline or of the non-noradrenergic motor transmitter (ATP), or both, has not been determined.

Blood vessels

The physiological significance of the inhibition by acetylcholine of noradrenergic transmission to blood vessels is uncertain. The vasodilator action of acetylcholine is mediated by the release of an endothelium-derived relaxing factor (EDRF) (Furchgott & Zawadski, 1980; Furchgott, 1983). However, the inhibitory effect of acetylcholine on noradrenergic transmission is not affected by removal of the endothelium, and therefore does not involve EDRF (Loiacono & Story, 1984). In some blood vessels, EDRF is not involved in the vasodilator response to acetylcholine (Brayden & Large, 1986).

Where there is a cholinergic innervation, as is the case for blood vessels of skeletal muscle (Burn, 1938; Uvnas, 1960, 1967), the vasodilator responses elicited by stimulation of these nerves could be due to inhibition of a noradrenergically maintained vasoconstrictor tone, as suggested by Vanhoutte & Shepherd (1973). However, there is little or no evidence for the existence of a cholinergic innervation in most of the blood vessels in which the inhibitory effect of acetylcholine on noradrenergic transmission has been demonstrated. There is some evidence for a cholinergic innervation of the blood vessels in the rabbit ear (Armin, Grant, Thompson & Tickner, 1953; Holton & Rand, 1962), and cholinesterases are associated with the periarterial sympathetic nerve fibres of the artery (Waterson et al., 1970). However, according to Hume & Waterson (1978), there are no cholinergic nerves associated with the rabbit ear artery. Atropine has only slight effects on the vasoconstrictor responses to sympathetic nerve stimulation in the rabbit ear (Holton & Rand, 1962) or the isolated artery (Rand & Varma, 1970), whereas blockade of an inhibitory component due to simultaneous release of acetylcholine would be expected to result in a greater vasoconstrictor response.

Blood vessels with a cholinergic innervation in which endogenously released acetylcholine has been shown to inhibit noradrenaline release include the canine coronary artery (Cohen, Shepherd & Vanhoutte, 1984) and gastric artery (Van Hee & Vanhoutte, 1978). Similar findings have been obtained in the perfused vasculature of the rabbit lung (Mathé et al. 1977).



Conclusion

There is overwhelming evidence for the presence of cholinoceptors on sympathetic noradrenergic nerve terminals and their distribution appears to be ubiquitous in terms of mammalian species and tissues innervated.

Activation of prejunctional nicotinic cholinoceptors results in release of noradrenaline and sometimes in facilitation of stimulation-induced noradrenaline release. Their physiological role is uncertain and the only attempt to assign them a role in the cholinergic link hypothesis of noradrenergic transmission has faded from prominence.

Activation of prejunctional muscarinic cholinoceptors results in inhibition of noradrenaline release and they appear to be involved in the physiological control of neurotransmission, at least in those tissues that contain a cholinergic as well as a noradrenergic innervation.

Other effects on noradrenergic transmission produced by cholinoceptor agonists are of uncertain significance and remain as curiosities.



Cardiac preparations:

Burn & Dutta, 1948 Heymans & Bennati, 1949 Jacob, 1952 Kottegoda, 1953a Szentiványi & Kövér, 1956 Holtz & Westermann, 1955 Burn & Rand, 1958a

Vascular preparations

Kottegoda, 1953b Burn & Rand, 1957, 1958b Burn, Leach, Rand & Thompson, 1959 Steinsland & Furchgott, 1975a,b

Spleen

Chernigovskii, 1940 Brandon & Rand, 1961 Daly & Scott, 1961 Macmillan & Rand, 1962 Boatman & Brody, 1964

Respiratory tract

Hawkins & Paton, 1958

Intestinal preparations

Ambache & Edwards, 1951 Gillespie & MacKenna, 1960, 1961.

Skin (piloerector muscles)

Brücke, 1935 Coon & Rothman, 1940 Burn et al., 1959 Burn & Rand, 1960a

Nictitating membrane

Thompson, 1958 Burn et al., 1959



```
2504075650
```

from (3H)-noradrenaline-labelled tissues by nicotinic agonists. See text for further references. Cardiac preparations Rabbit Richardson & Woods, 1959 Lindmar & Muscholl, 1961 Lindmar, 1962 Muscholl & Maître, 1963 Lindmar et al., 1967, 1968 Löffelholz, 1967, 1970a,b Löffelholz & Muscholl, 1969 Fozard & Muschell, 1972 Muscholl, 1973b Göthert, 1974 Starke & Montel, 1974 Dubey, Muscholl & Pfeiffer, 1975 Göthert, Kennerknecht & Thielecke, 1976 Sarantos-Laska, Majewski, McCulloch & Rand, 1980a Guinea-pig Bhagat, Robinson & West, 1967 Lindmar et al., 1968 Westfall, 1971a Westfall & Brasted, 1972, 1973, 1974 Allen, Glover, Rand & Story, 1972b Sorimachi, Oesch & Thoenen, 1973 Westfall & Hunter, 1974. Rat Sharma & Banerjee, 1978 Fuder, Siebenborn & Muscholl, 1982b (but see Westfall & Saunders, 1977) Spleen (cat) Brandon & Boyd, 1961 Blakeley, Brown & Ferry, 1963 Herrting & Widholm, 1965 Fischer, Weise & Kopin, 1966 Davey, Haydon & Scholfield, 1968 Krauss, Carpenter & Kopin, 1970 Kirpekar, Prat, Puig & Wakade, 1972 Vascular preparations (rabbit) Aorta Kiran & Khairallah, 1969 Pulmonary artery Su & Bevan, 1970 Nedergaard & Schrold, 1973, 1977 Starke & Weitzell, 1978 Ear artery Allen, Glover, McCulloch, Rand & Story, 1975 Intestinal tissue Rabbit colon Gillespie & MacKenna, 1960 Guinea-pig taenia caeci Kuchii, Miyahara & Shibata, 1973 Vas deferens (rat)

Table 2. Reports of release of noradrenaline or of radioactivity



Jayasundar & Vohra, 1978

Table 3. Evidence for a cholinergic component in the predominantly noradrenergic sympathetic nerve supply. See text for further references.

Cat

Nictitating membrane

Bacq & Fredericq, 1935 Burn & Rand, 1960a Nystrom, 1962

Heart

Folkow, Frost, Halger & Uvnas, 1948 Day & Rand, 1961

Spleen

Burn & Rand, 1960a Brandon & Rand, 1961

Uterus

Sherif, 1935 Burn & Rand, 1960a

Hind leg

Folkow et al.,1948

Dog

Hind leg

Bülbring & Burn, 1935 Bogaert, De Schaepdryver & De Vleeschhouwer, 1961 McCubbin, Kaneko & Page, 1961

Tongue

von Euler & Gaddum, 1931

Rabbit

Intestine

Gillespie & MacKenna, 1959 Day & Rand, 1961

Atria

Hukovic, 1959

Ear

Burn & Rand, 1960a Holton & Rand, 1962



References

- Allen, G.S., Glover, A.B., McCulloch, M.W., Rand, M.J. & Story, D.F. (1975) Modulation by acetylchcline of adrenergic transmission in the rabbit ear artery. Br. J. Pharmacol., 54, 49-53.
- Allen, G.S., Glover, A.B., Rand, M.J. & Story, D.F. (1972b) Effects of acetylcholine on vasoconstriction and release of ¹H-noradrenaline in response to sympathetic nerve stimulation in the isolated artery of the rabbit. Br. J. Pharmacol., 46, 527-528P.
- 3 Allen, G.S., Rand, M.J. & Story, D.F. (1972a) Effects of MoN-A-343 and DMFP on the uptake and release of ³H-noradrenaline by guinea-pig atria. Br. J. Pharmacol., 45, 480-489.
- Allen, G.S., Rand, M.J. & Story, D.F. (1973b) Comparison of the effects of six cholinomimetic drugs on inhibition of uptake of 3H-(±)-noradrenaline by guinea-pig atria. Br. J. Pharmacol., 47, 179-180.
- 5 Allen, G.S., Rand, M.J. & Story, D.F. (1974) Effects of the muscarinic agonist McN-A-343 on release by sympathetic nerve stimulation of 3H-noradrenaline from rabbit isolated ear arteries and guinea-pig atria. Br. J. Pharmacol., 51, 29-34.
- 6 Ambache, N. & Edwards, J. (1951) Reversal of nicotine action on the intestine by atropine. Br. J. Pharmacol., 6, 311-317.
- Armin, J., Grant, R.T., Thompson, R.H.S and Tickner, A. (1953) An explanation for the heightened vascular reactivity of the denervated rabbit's ear. J. Physiol., 121, 603-622.
- 8 Armitage, J.A. & Burn, J.H. (1967) Effect of physostigmine on the contraction of the retractor penis in the dog in response to sympathetic stimulation. Br. J. Pharmacol., 29, 218-229.
- Bacq, Z.M. & Fredericq, H. (1935) Essai d'identification du médiateur chimique libéré dans la membrane nictitante du chat par l'excitation sympathique. Arch. int. Physiol., 40, 297-310.
- 10 Baldessarini, R.J. (1975) Release of catecholamines. In Handbook of Psychopharmacology, Vol.3, (Eds L.L.Iversen, S.D. Iversen and S.H.Snyder) pp.37-137, Plenum Press, New York.
- 11 Bell, C. (1968) Differential effects of tetrodotoxin on sympathomimetic actions of nicotine and tyramine. Br. J. Pharmacol., 32, 96-103.
- 12 Bentley, G.A. (1962) Studies on sympathetic mechanisms in isolated intestinal and vas deferens preparations. Br. J. Pharmacol., 19, 85-98.
- Bernard, P.J. & De Schaepdryver, A.F. (1964) Adrenergic mechanisms in the dog hind leg. Arch. int. Pharmacodyn., 148, 301-305.
- 14 Bevan, J.A. & Hauesler, G. (1975) Electrical events associated with the action of nicotine at the adrenergic nerve terminal. Arch. int. Pharmacodyn., 218, 84-95.
- Bevan, J.A. & Su, C. (1964) The sympathetic mechanism in the isolated pulmonary artery of the rabbit. Br. J. Pharmacol., 22, 176-182.
- 16 Bevan, J.A. & Su, C. (1972) Uptake of nicotine by the sympathetic nerve terminals in the blood vessel. J. Pharmacol. Exp. Ther., 182, 419-426.
- 17 Bhagat, B. (1966) Response of isolated guinea-pig atria to various ganglion-stimulating agents. J. Pharmacol. Exp. Ther., 154, 264-270.
- Bhagat, B., Robinson, I.M. & West, W.L. (1967) Mechanism of sympathomimetic responses of isolated guinea-pig atria to nicotine and dimethylphenylpiperazinium iodide. Br. J. Pharmacol., 30, 470-477.



- 19 Birmingham, A.T. & Wilson, A.B. (1965) Analysis of the blocking action of dimethyphenylpiperazinium iodide on the inhibition of isolated small intestine produced by stimulation of the sympathetic nerves. Br. J. Pharmacol., 24, 375-386.
- 20 Blakeley, A.G.H., Brown, G.L. & Ferry, C.B. (1963)
 Pharmacological experiments on the release of the sympathetic transmitter. J. Physiol., 167, 504-514.
- 21 Boatman, D.L. & Brody, M.J. (1964) Analysis of vascular responses of the spleen. Am. J. Physiol., 207, 155-161.
- 22 Boatman, D.L., Shaffer, R.A., Dixon, R.L. & Brody, M.J. (1965) Function of vascular smooth muscle and its sympathetic innervation in the new born dog. J. Clin. Invest., 44, 241-246.
- 23 Bogaert, M., De Schaepdryver, A.F. & De Vleeschhouwer, G.R. (1961) On the pharmacology of guanethidine. Arch. int. Pharmacodyn., 34, 224-236.
- 24 Boura, A.L.A. & Green, A.F. (1959) The actions of bretylium: adrenergic neurone blocking and other effects. Br. J. Pharmaccl., 14, 536-548.
- Bowman, W.C., Callingham, B.A. & Cuthbert, A.W. (1964) The effects of physostigmine on the mechanical and electrical responses of the cat nictitating membrane. Br. J. Pharmacol., 22, 558-576.
- 26 Boyd, H., Chang, V. & Rand, M.J. (1960) The anticholinesterase activity of some antiadrenaline drugs. Br. J. Pharmacol., 15, 525-531.
- 27 Boyd, H., Chang, V. & Rand, M.J. (1961) The local anaesthetic activity of bretylium in relation to its action in blocking sympathetic responses. Arch. int. Pharmacodyn., 131, 10-23.
- 28 Brandon, K.W. & Boyd, H. (1961) Release of noradrenaline from the spleen of the cat by acetylcholine. Nature, 192, 880-881.
- 29 Brandon, K.W. & Rand, M.J. (1961) Acetylcholine and the sympathetic innervation of the spleen. J. Physiol., 157, 18-32.
- 30 Brayden, J.E. & Large, W.A. (1986) Electrophysiological analysis of neurogenic vasodilatation in the isolated lingual artery of the rabbit. Br. J. Pharmacol., 89, 163-171.
- Briley, M.S., Langer, S.Z. & Story, D.F. (1979) Binding studies on alpha-adrenoceptors and muscarinic cholinoceptors in rat heart ventricle: effect of chemical sympathectomy. Br. J. Pharmacol., 66, 90P.
- 32 Brücke, F.T. (1935) Über die Wirkung von Acetylcholin auf die Pilomotoren. Klinische Wochenschrift, 14, 7-9.
- 33 Bülbring, E. & Burn, J.H. (1935) The sympathetic dilator fibres in the muscles of the cat and dog. J. Physiol, 83, 483-501.
- 34 Bunge, R., Johnson, M. & Rees, C. (1978) Nature and nurture in the development of the autonomic neurone. Science, 199, 1409-1416.
- 35 Burn, J.H. (1938) Sympathetic vasodilator fibres. Physiol. Rev., 18, 137-153.
- 36 Burn, J.H. (1961) A new view of adrenergic nerve fibres, explaining the actions of reserpine, bretylium and guanethidine. Br. Med. J., 1, 1623-1627.
- 37 Burn, J.H. (1967) Release of noradrenaline from the sympathetic postganglionic fibre. Br. Med. J., 2, 197-201.
- 38 Burn, J.H. (1968) The development of the adrenergic fibres. Br. J. Pharmacol., 32, 575-582.
- 39 Burn, J.H. (1971) Release of noradrenaline from sympathetic nerve endings. Nature, 231, 237-240.



- 40 Burn, J.H. (1977a) Evidence that acetylcholine releases noradrenaline in the sympathetic fibre. J. Pharm. Pharmacol., 29, 325-329.
- Burn, J.H. (1977b) The function of acetylcholine released from sympathetic fibres. Clin. Exp. Pharmacol. Physiol., 4, 59-100.
- Burn, J.H. & Dutta, N.K. (1948) The action of antagonists of acetylcholine on the vessels of the rabbit ear. Br. J. Pharmacol., 3, 354-361.
- Burn, J.H. & Gibbons, W.R. (1964) The sympathetic postganglionic fibre and the block by bretylium: the block prevented by hexamethonium and imitated by mecamylamine. Br. J. Pharmacol., 22, 549-557.
- Burn, J.H., Leach, E.H., Rand, M.J. & Thompson, J.W. (1959)
 Peripheral effects of nicotine and acetylcholine resembling those of sympathetic nerve stimulation. J. Physiol., 148, 332-352.
- 45 Burn, J.H. & Rand, M.J. (1957) Reserpine and noradrenaline in artery wall. Lancet, 273, 1097.
- 46 Burn, J.H. & Rand, M.J. (1958a) Action of nicotine on the heart. Br. Med. J., 1,137-139.
- 47 Burn, J.H. & Rand, M.J. (1958b) Noradrenaline in artery walls and its dispersal by reserpine. Br. Med. J., i, 903-908.
- 48 Burn, J.H. & Rand, M.J. (1959) Sympathetic postganglionic mechanism. Nature, 184, 163-165.
- 49 Burn, J.H. & Rand, M.J. (1960a) Sympathetic postganglionic cholinergic fibres. Br. J. Pharmacol., 15, 56-66.
- 50 Burn, J.H. & Rand, M.J. (1960b) New observations on the sympathetic postganglionic mechanism. Am. J. Med., 29, 1002-1007.
- 51 Burn, J.H. & Rand, M.J. (1962) A new interpretation of the adrenergic nerve fiber. Adv. Pharmacol. Chemother., 1, 1-30.
- 52 Burn, J.H. & Rand, M.J. (1965) Acetylcholine in adrenergic transmission. Annu. Rev. Pharmacol., 5, 163-182.
- 53 Burn, J.H., Rand, M.J. & Wein, R. (1963) The adrenergic mechanism in the nictitating membrane. Br. J. Pharmacol., 20, 83-94.
- 54 Burn, J.H. & Weetman, D.F. (1963) The effect of eserine on the response of the vas deferens to hypogastric nerve stimulation. Br. J. Pharmacol., 20, 74-81.
- 55 Burnstock, G., Campbell, G. & Rand, M.J. (1966) The inhibitory innervation of the taenia of the guinea-pig caecum. J. Physiol., 182, 504-526.
- 56 Cabrera, R., Torrance, R.W. & Viveros, H. (1966) The action of acetylcholine and other drugs upon the terminal parts of the postganglionic sympathetic fibre. Br. J. Pharmacol., 27, 51-63.
- 57 Campbell, G. (1970) Autonomic nervous supply to effector tissues. In Smooth Muscle (Eds E. Bulbring, A.F. Brading, A.W. Jones and T.Tomita) Arnold, London, pp 451-495.
- 58 Chernigovskii, V.N. (1940) Issledovanie retseptorov nekotorykh vnutrennikh organov. Soobshchenie III. Deïstvie atsetilkholina, nikotina, histamina i KCl na retseptory selezenki. Fiziol. Zh. SSSR, 29, 526-535.
- 59 Choo, L.K., Mitchelson and Vong, Y.M. (1985) The interaction of McN-A-343 with pirenzepine and other selective muscarinic antagonists at a prejunctional muscarinic receptor. Naunyn-Schmiedeberg's Arch. Pharmacol., 328, 430-438.



- 60 Choo, L.K., Mitchelson, F. & Vong, Y.M. (1986) Interaction of some muscarinic agonists and antagonists at the prejunctional muscarinic receptor in the rabbit ear artery preparation. Br. J. Pharmacol., 87, 733-740.
- 61 Cohen, R.R., Shepherd, J.T. & Vanhoutte, P.M. (1984)
 Neurogenic cholinergic prejunctional inhibition of sympathetic
 beta adrenergic relaxation in the canine coronary artery. J.
 Pharmacol. Exp. Ther., 229, 417-421.
- 62 Collett, A.R. & Story, D.F. (1984) Effects of adrenoceptor antagonists and neuronal uptake inhibitors on dimethylphenylpiperazinium-induced release of catecholamines from the rabbit isolated adrenal gland and guinea-pig atria. J. Pharmacol. Exp. Ther., 231 379-386.
- 63 Comer, M.S. & Di Palma, J.R. (1961) TM10 activity in cholinergic compounds. Arch. int. Pharmacodyn., 131, 368-377.
- 64 Concha, J. & Norris, B. (1968) Studies on renal vasomotion. Br. J. Pharmacol. 34, 277-290.
- 65 Coon, J.M. & Rothman, S. (1940) Nature of pilomotor response to acetylcholine, some observations on pharmaco-dynamics of skin. J. Pharmacol. Exp. Ther., 68, 301-311.
- 66 Cooper, T. (1965) In Nervous Control of the Heart (Ed. Randall, W.C.) Williams & Wilkins, Baltimore, p 130.
- 67 Dale, H.H. & Feldberg, W. (1934) The chemical transmission of secretory impulses to the sweat glands of the cat. J. Physiol, 82, 121-128.
- 68 Daly, M. de B. & Scott, M.J. (1961) The effects of acetylcholine on the volume and vascular resistance of the dog's spleen. J.Physiol, 156, 246-259.
- 69 Davey, M.J., Hayden, M.L. & Scholfield, P.C. (1968) The effects of bretylium on C fibre excitation and noradrenaline release by acetylcholine and electrical stimulation. Br. J. Pharmacol., 34, 377-387.
- 70 Day, M.D. & Rand, M.J. (1961) Effect of guanethidine in revealing cholinergic sympathetic fibres. Br. J. Pharmacol., 17, 245-260.
- 71 de la Lande, I.S. & Rand, M.J. (1965) A simple isolated nerveblood vessel preparation. Austral. J. exp. Biol. & Med. Sci., 43, 639-656.
- 72 Dixon, W.E. (1924) Nicotin, Piperidin, Lupetidin, Ceytisin, Lobelin, Spartein, Gelsemin. Mittel, welche auf bestimmte Nervenzellen Wirken. Heffter-Haubner's Handbuch experimentellen Pharmakologie, 2, 656-736.
- 73 Dubey, M.P., Muscholl, E. & Pfeiffer, A. (1975) Muscarinic inhibition of potassium induced noradrenaline release and its dependence on the calcium concentration. Naunyn-Schmiedeberg's Arch. Pharmacol., 291, 1-15.
- 74 Edvinsson, L., Falck, B. & Owman, C. (1977) Possibilities for a cholinergic action on smooth musculature and on sympathetic axons in brain vessels mediated by muscarinic and nicotinic receptors. J. Pharmacol. Exp. Ther., 200, 117-126.
- 75 Ehinger, B. & Falck, B. (1966) Concomitant adrenergic and parasympathetic fibres in the rat iris. Acta Physiol. Scand., 67, 201-207.
- 76 Ehinger, B., Falck, B., Persson, H., Rosengren and Sporrong, B. (1970) Acetylcholine in adrenergic terminals of the cat iris. J. Physiol., 209, 557-565.
- 77 Endoh, M. & Hashimoto, K. (1970) Pharmacological evidence of autonomic nerve activities in canine papillary muscle. Am. J. Physiol., 218, 1459-1463.

29100201073

- Fresynaptic receptor systems on the noradrenergic neurones of the rabbit pulmonary artery. Naunyn-Schmiedeberg's Arch. Pharmacol., 296, 229-247.
- 79 Endoh, M., Tamura, K. & Hashimoto, K. (1970) Negative and positive inotropic responses of the blood-perfused canine papillary muscle to acetylcholine. J. Pharmacol. Exp. Ther., 175, 377-387.
- 80 Engel, V. & Löffelholz, K. (1976) Presence of muscarinic inhibitory and absence of nicotinic excitatory receptors at the terminal sympathetic nerves of chicken hearts. Naunyn-Schmiedeberg's Arch. Pharmacol., 295, 225-230.
- Schmiedeberg's Arch. Pharmacol., 295, 225-230.

 81 Eränkö, O., Rechardt, L., Eränkö, L. & Cunningham, A. (1970)
 Light and electron microscope histological observations on
 cholinesterase-containing sympathetic nerve fibres in the
 pineal body of the rat. Histochem.J., 2, 479-489.
- 82 Esterhuizen, A.C., Graham, J.D.P., Lever, J.D. & Spriggs, T.L.B. (1968) Catecholamine and acetylcholinesterase distribution in relation to noradrenaline release. An enzyme histochemical and autoradiographic study on the innervation of the cat nictitating membrane. Br. J. Pharmacol., 32, 46-56.
- Ferry, C.B. (1963) The sympathetic effect of acetylcholine on the spleen of the cat. J. Physiol., 167, 487-504.
- 84 Ferry, C.B. (1966) Cholinergic link hypothesis in adrenergic neuroeffector transmission. Physiol. Rev., 46, 420-456.
- Fewings, J.D., Rand, M.J., Scroop, G.C. & Whelan, R.F. (1966)
 The action of nicotine on the blood vessels of the hand and forearm in man. Br. J. Pharmacol., 26, 567-579.
- Fischer, J.E., Weise, V.K. & Kopin, I.J. (1966) Interactions of bretylium and acetylcholine at sympathetic nerve endings. J. Pharmacol. Exp. Ther., 153, 523-529.
- 87 Folkow, B., Frost, J., Haeger, K. & Uvnas, B. (1948) Cholinergic fibres in sympathetic outflow to heart in dog and cat. Acta Physiol. Scand., 15, 421-426
- 88 Fozard, J. (1979) Cholinergic mechanisms in adrenergic function. In Trends in Autonomic Pharmacology (Ed S. Kalsner. pp 145-194. Urban & Schwarzenburg, Baltimore.
- 89 Fozard, J. & Muscholl, E. (1972) Effects of several muscarinic agonists on cardiac performance and the release of noradrenaline from sympathetic nerves of the perfused rabbit heart. Br. J. Pharmacol., 45, 616-629.
- 90 Fozard, J. & Muscholl, E. (1974) Atropine-resistant effects of the muscarinic agonists McN-A-343 and AHR 602 on cardiac performance and the release of noradrenaline from sympathetic nerves of the perfused rabbit heart. Br. J. Pharmacol., 50, 531-541.
- 91 Fozard, J. & Mwaluko, G.M.P. (1976) Mechanism of the indirect sympathomimetic effect of 5-hydroxytryptamine on the isolated heart of the rabbit. Br. J. Pharmacol., 57, 115-125.
- 92 Fuder, H. (1982) The affinity of pirenzepine and other antimuscarinic compounds for pre- and postsynaptic muscarinic receptors of the isolated rabbit and cat heart. Scand. J. Gastroenterol., 17 (Suppl.72), 79-85.
- 93 Fuder, H., Meiser, C., Wormstall, H. & Muscholl, E. (1981) The effects of several muscarinic antagonists on pre- and postsynaptic receptors in the isolated rabbit heart. Naunyn-Schmiedeberg's Arch. Pharmacol., 316, 31-37.
- 94 Fuder, H. & Muscholl, E. (1974) The effect of methacholine on noradrenaline release from the rabbit heart perfused with indomethacin. Naunyn-Schmiedeberg's Arch. Pharmacol., 285, 127-132.



- 95 Fuder, H., Rink, D. & Muscholl, E. (1982a) Sympathetic nerve stimulation on the perfused rat heart. Affinities of N-methylatropine and pirenzipine at pre and postsynaptic muscarine receptors. Naunyn-Schmiedeberg's Arch. Pharmacol., 318, 210-219.
- 96 Fuder, H., Schimkus, R. & Braun, H.J. (1985) Affinity and efficacy of selected agonists in the a2-adrenoceptor mediated inhibition of 3H-noradrenaline release. In Pharmacology of Adrenoceptors (Ed E. Szabadi, C.M. Bradshaw and R. Nahorski), p 335. Macmillan Press, London.
- 97 Fuder, H., Siebenborn, R. & Muscholl, E. (1982b) Nicotine receptors do not modulate the ³H-noradrenaline release from the isolated heart evoked by sympathetic nerve stimulation. Naunyn-Schmiedeberg's Arch. Pharmacol., 381, 301-307.
- 98 Furchgott, R.F. (1983) Role of endothelium in responses of vascular smooth muscle. Circ. Res., 53, 557-573.
- 99 Furchgott, R.F., Steinsland, O.S. & Wakade, T.D. (1975)
 Studies on prejunctional muscarinic and nicotinic receptors.
 In Chemical Tools in Catecholamine Research (Eds. O. Almgren,
 A. Carlsson & J. Engel) North Holland Amsterdam, pp 167-174.
- 100 Furchgott, R.F. & Zawadski, J.V. (1980) The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. Nature, 288, 373-376.
- 101 Garcia-Seville, J.A., Dubocovich, M.L. & Langer, S.Z. (1985) Interaction between presynaptic facilitatory angiotensin II receptors and autoinhibitory muscarinic cholinoceptors on ³H-noradrenaline release in the rabbit heart. Naunyn-Schmiedeberg's Arch. Pharmacol., 330, 9-15.
- 102 Gardiner, J.E., Hellmann, K. & Thompson, J.W. (1962) The nature of the innervation of the smooth muscle, Harderian gland and blood vessels of the cat's nictitating membrane. J. Physiol.. 163, 436-456.
- Physiol., 163, 436-456.

 103 Gillespie, J.S. & MacKenna, B.R. (1959) The effect of reserpine of the rabbit ileum and colon to stimulation of the extrinsic nerves in vitro. J. Physiol, 147, 31P-32P.
- 104 Gillespie, J.S. & MacKenna, B.R. (1960) The inhibitory action of nicotine on the rabbit colon. J. Physiol., 152, 191-205.
- 105 Gillespie, J.S. & MacKenna, B.R. (1961) The inhibitory action of the sympathetic nerves on the smooth muscle of the rabbit gut, its reversal by reserpine and restoration by catechol amines and DOPA. J. Physiol., 156, 17-34.
- 106 Ginzel, K.H. & Kottegoda, S.R. (1953) Nicotine-like actions in auricles and blood vessels after denervation. Br. J. Pharmacol., 8, 348-351.
- 107 Göthert, M. (1974) Effects of halothane on the sympathetic nerve terminals of the rabbit heart. Differences in membrane actions of halothane and tetracaine. Naunyn-Schmiedeberg's Arch. Pharmacol., 286,125-143.
- 108 Göthert, M. (1977) Effects of presynaptic modulators on Ca²+induced noradrenaline release from cardiac sympathetic nerves.
 Naunyn-Schmiedeberg's Arch. Pharmacol., 300, 267-272.
- 109 Göthert, M., Kennerknecht, E. & Thielecke, G. (1976)
 Inhibition of receptor-mediated noradrenaline release from the sympathetic nerves of the isolated rabbit heart by anaesthetics and alcohols in proportion to their hydrophobic property. Naunyn-Schmiedeberg's Arch. Pharmacol., 292, 145-142.



- 110 Graham, J.D.P., Lever, J.D. & Spriggs, T.L.B (1968) An examination of adrenergic axons around pancreatic arterioles of the cat for the presence of acetylcholinesterase by high resolution autoradiographic and histochemical methods. Br. J.
- Pharmacol., 33, 15-20.
 111 Gulati, O.D. & Panchal, D.I. (1978) Some observations on the development of adrenergic innervation in rabbit intestine. Br. J. Pharmacol., 64, 247-251.
- 112 Haeusler, G., Haefely, W. & Huerlimann, A. (1969a) On the mechanism of the adrenergic nerve blocking action of bretylium. Naunyn-Schmiedeberg's Arch. Pharmacol., 265, 260-
- 113 Haeusler, G., Thoenen, H., Haefely, W. & Huerlimann, A. (1968) Electrical events in cardiac adrenergic nerves and noradrenaline release from the heart induced by acetylcholine and KCl. Naunyn-Schmiedeberg's Arch. Pharmacol., 261, 389-411.
- 114 Haeusler, G., Thoenen, H., Haefely, W. & Huerlimann, A. (1969b) Electrosekretorische Koppelung bei der Noradrenalin-Freisetzung aus adrenergen Nevernfasen durch nicotinartig wirkende Substanzen. Naunyn-Schmiedeberg's Arch. Pharmacol., 263, 217-218.
- 115 Haney, H.F. & Lindgren, A.J. (1945) The effect of acetylcholine on the atropinized denervated heart. Physiol., 145, 177-180.
- 116 Hawkins, D.F. & Paton, W.D.M. (1958) Responses of isolated bronchial muscle to ganglionically active drugs. J. Physiol., 144, 193-219.
- 117 Hellmann, K. (1963) The isolated pilomotor muscle as an in J. Physiol., 123, 289-300. vitro preparation.
- 118 Herrting, G. & Widholm, S. (1965) Über den Mechanismus der Noradrenalin-Freisetzung aus sympathischen Nervenendigungen. Naunyn-Schmiedberg's Arch. Exp. Path. Pharmacol., 250, 257-258.
- 119 Heymans, C. & Bennati, D. (1949) Méchanisme de la tachycardie
- par l'acétylcholine. Arch. int. Pharmacodyn., 79, 486-489.
 120 Higgins, C.B., Vatner, S.F. & Braunwald, E. (1973)
 Parasympathetic control of the heart. Pharmacol. Rev., 25, 119.
- 121 Hobbinger, F., Mitchelson, F. & Rand, M.J. (1969) The actions of some cholinomimetic drugs on the isolated taenia of the guinea-pig caecum. Br. J. Pharmacol., 36, 53-69.
- 122 Hoffmann, F., Hoffmann, E.J., Middleton, S. & Talesnik, J. (1945) The stimulating effect of acetylcholine on the mammalian heart and the liberation of an epinephrine-like substance by the isolated heart. Am. J. Physiol., 144, 189-198.
- 123 Hökfelt, T. (1967) Ultrastructural studies on adrenergic nerve terminals in the albino rat iris after pharmacological and experimental treatment. Acta Physiol.Scand., 69, 125-126.
- 124 Hollenberg, M., Carriere, S. & Barger, A.C. (1965) Biphasic action of acetylcholine on ventriocular myocardium. Res., 16, 527-536.
- 125 Holton, P. & Rand, M.J. (1962) Sympathetic vasodilatation in the rabbit ear. Br. J. Pharmacol., 19, 513-526.
- 126 Holtz, P. & Westermann, E. (1955) Versuche mit Acetyl-, Propionyl- und Butyryl-cholin am isolierten Herzvorhofpräparat. Arch. exp. Pathol. Pharmakol., 225, 421-427.



- 127 Hope, W., McCulloch, M.W., Rand, M.J. & Story, D.F. (1974) Cholinergic effects on adrenergic transmission in rabbit and guinea-pig isolated atria. Proc. Aust. Physiol. Pharmacol. Soc., 5, 213-314.
- 128 Hope, W., McCulloch, M.W., Rand, M.J. & Story, D.F. (1978) The effect of calcium on the interaction between acetylcholine and noradrenergic transmission in the rabbit ear artery. Clin. Exp. Pharmacol. Physiol., 5, 290.
- 129 Hukovic, S. (1959) Isolated rabbit atria with sympathetic nerve supply. Br. J. Pharmacol., 14, 372-376.
- 130 Hukovic, S. (1960) The action of sympathetic blocking agents on isolated and innervated atria and vessels. Br. J. Fharmacol., 15, 117-121.
- 131 Hukovic, S. (1966) The effect of anticholinestreases on the increase in rate of the isolated heart in response to sympathetic stimulation. Br. J. Pharmacol., 28, 273.
- 132 Hume, W.R., de la Lande, I.S. & Waterson, J.G. (1972) Effect of acetylcholine on the response of the isolated rabbit ear artery to stimulation of the perivascular sympathetic nerves. Eur. J. Pharmacol., 17, 227-233.
- 133 Hume, W.R. & Waterson, J.G. (1978) The innervation of the rabbit ear artery. An ultrastructural study. Blood Vessels, 15, 348-364.
- 134 Iwayama, T., Furness, J.B. & Burnstock, G. (1970) Dual adrenergic and cholinergic innervation of the cerebral arteries of the rat, an ultrastructural study. Circ. Res., 26, 635-646.
- 135 Jacob, J. (1952) Antagonistes de la stimulation acétylcholinique du coeur isolé et atropinisé de lapin comparison entre divers agents synaptolytiques, antihistaminiques et antifibrillants. Arch. int. Pharmacodyn.. 89. 106-112.
- Pharmacodyn., 89, 106-112.

 136 Jacobowitz, D. & Koelle, G.B. (1965) Histochemical correlations of acetylcholinesterase and catecholamines in postganglionic autonomic nerves of the cat, rabbit and guinea-pig. J. Pharmacol. Exp. Ther., 148, 225-237.
- 137 Jaju, B.D. (1969) Burn and Rand's hypothesis. Indian J.Physiol.Pharmacol. 13, 1-27.
- 138 Jayasundar, S. & Vohra, M.M. (1978) An analysis of action of nicotinic agents on adrenergic nerve terminals in rat isolated vas deferens. Arch. int. Pharmacodyn., 232, 192-201.
- 139 Jones, M.E.L. & Spriggs, T.L.B. (1975) An inhibitory effect of atropine on responses of the vas deferens of the mouse to field stimulation. Br. J. Pharmacol., 54, 339-349.
- 140 Kiran, B.K. & Khairallah, P.A. (1969) Angiotensin and norepinephrine efflux. Eur. J. Pharmacol., 6, 102-108.
- 141 Kirpekar, S.M., Garcia, A.G. & Prat, J.C. (1980) Action of nicotine on sympathetic nerve terminals. J. Pharmacol. Exp. Ther., 213, 133-138.
- 142 Kirpekar, S.M., Prat, J.C., Puig, M. & Wakade, A.R. (1972) Modification of the evoked release of noradrenaline from the perfused cat spleen by various ions and agents. J. Physiol., 221, 601-615.
- 143 Kirpekar, S.M., Prat, J.C. & Wakade, A.R. (1975) Effect of calcium on the relationship between frequency of stimulation and release of noradrenaline from the perfused spleen of the cat. Naunyn-Schmiedeberg's Arch. Pharmacol., 287, 205-212.
- 144 Klinge, E. & Sjöstrand, N.O. (1977) Suppression of the excitatory adrenergic neurotransmission, a possible role of cholinergic nerves in the retractor penis muscle. Acta Physiol. Scand., 100, 368-376.



- 145 Knoll, J., Somogyi, G.T., Illes, P. & Vizi, E.S. (1972)
 Acetylcholine released from isolated vas deferens of the rat.
 Naunyn-Schmiedeberg's Arch. Pharmacol., 274, 198-202.
- 146 Kosterlitz, H.W. & Lees, G.M. (1972) Interrelationships between adrenergic and cholinergic mechanisms. In Catecholamines. Handbook of Experimental Pharmacology, Vol 33, (Ed H. Blaschko and E Muscholl), pp.762-812, Springer, Berlin, Heidelberg, New York.
- 147 Kottegoda, S.R. (1953a) Stimulation of isolated rabbit auricles by substances which stimulate ganglia. Br. J. Pharmacol.. 8, 83-86.
- Pharmacol., 8, 83-86.

 148 Kottegoda, S.R. (1953b) The action of nicotine and acetylcholine on the vessels of the rabbit's ear. Br. J. Pharmacol., 8, 156-161.
- 149 Krauss, K.R., Carpenter, D.O. & Kopin, I.J. (1970)

 Acetylcholine-induced release of norepinephrine in the presence of tetrodotoxin. J. Pharmacol. Exp. Ther., 173, 416-421.
- 150 Kuchii, M., Miyahara, J.T. & Shibata, S. (1973) (3H)-adenine nucleotide and (3H)-noradrenaline release evoked by electrical field stimulation, perivascular nerve stimulation and nicctine from the taenia of the guinea-pig caecum. Br. J. Pharmacol., 49, 258-267.
- 151 Kukovetz, W.R. (1962) Kontractilität und Phosphorylaseaktivität des Herzens bei ganglionärer Erregung nach adrenerger Blockade and unter Atropin. Arch. exp. Pathol. Pharmakol., 243, 391-406.
- 152 Lamble, A.P. & Lamble, J.W. (1975) Some effects of pilocarpine and acetylcholine on the responses of the guinea-pig vas deferens to hypogastric nerve stimulation and transmural stimulation and applied 1-noradrenaline. J. Pharm. Pharmacol., 27, 79-87.
- 153 Langley, A.E. & Gardier, R.W. (1974) Regulation of norepinephrine and dopamine-β-hydroxylase release during sympathetic nerve stimulation. Fed. Proc., 33, 523 Abs.
- sympathetic nerve stimulation. Fed. Proc., 33, 523 Abs.
 154 Langley, A.E. & Gardier, R.W. (1977) Effect of atropine and acetylcholine on nerve stimulated output of noradrenaline and dopamine-beta-hydroxylase from isolated rabbit and guinea-pig hearts. Naunyn-Schmiedeberg's Arch. Pharmacol., 297, 251-256.
- 155 Lauer, J.A. & Steinsland, O.S. (1983) Pirenzepine has similar affinities for pre- and post-junctional muscarinic receptors. Fed. Proc., 42, 1145.
- 156 Lavallée, M., de Champlain, C., Nadeau, A. & Yamaguchi, N. (1978) Muscarinic inhibition of endogenous myocardial catecholamine liberation in the dog. Can. J. Physiol. Pharmacol., 56, 642-649.
- 157 Leach, G.D.H. & Zumani, E.C. (1969) The effects of some changes in the perfusion solution on the vasoconstrictor responses of the isolated rat mesentery preparation. Br. J. Pharmacol., 36, 209-210.
- 158 Lederer, E., Rand, M.J. & Wilson, J. (1970) Evidence for the participation of a cholinergic receptor in adrenergic transmission. Arch. int. Pharmacodyn., 185, 105-120.
- 159 Lee, W.C. & Shideman, F.E. (1959) Mechanism of the positive inotropic response to certain ganglionic stimulants. J. Pharmacol. Exp. Ther., 126, 239-249.
- 160 Leighton, H.J. & Westfall, T.C. (1976) Role of impulse flow in the muscarinic inhibitory control of ³H-norepinephrine release from guinea-pig vasa deferentia. Fed. Proc., 35, 406.

Z9100201078

- 161 Leung, E. & Mitchelson, F. (1983) Muscarinic receptor inhibition and other effects of pancuronium in the rabbit ear artery preparation. Clin. Exp. Pharmacol. Physiol., 10, 629-639
- 171 Levy, M.N. (1971) Sympathetic-parasympathetic interactions in the heart. Circ. Res., 29, 437-445.
- 172 Levy, M.N. (1984) Cardiac sympathetic-parasympathetic
- interactions. Fed. Proc., 43, 2598-2602 173 Levy, M.N. & Blattberg, B. (1976) Effect of vagal stimulation on the overflow of norepinephrine into the coronary sinus during cardiac sympathetic nerve stimulation in the dog. Circ. Res., 38, 81-85.
- 174 Levy, M.N. & Martin, P.J. (1979) Neural control of the heart. In Handbook of Physiology. Cardiovascular System, The Heart. (Ed R.M. Berne, N.Sperelakis and S.R.Geiger), pp 581-620. Bethesda, Am. Physiol. Soc.
- 175 Levy, M.N., Ng, M., Martin, P. & Zieske, H. (1966) Sympathetic and parasympathetic interactions upon the left ventricle of the dog. Circ. Res., 19, 5-10.
- 176 Levy, M.N. & Zieske, H. (1969) Autonomic control of cardiac pacemaker activity and atrioventricular transmission. J. Appl. Physiol., 27, 465-470.
- 177 Li, C.K. & Mitchelson, F. (1980) The selective antimuscarinic action of stercuronium. Br. J. Pharmacol., 70, 313-321.
- 178 Lindamood, C., Johnson, S. & Fleming, W.W. (1978) Dual excitatory effect of acetylcholine in the mouse vas deferens. Proc. Soc. Exp. Biol. Med., 157, 200-201.
- 179 Lindmar, R. (1962) Die Wirkung von 1,1-Dimethylphenylpiperazinium Iodid am isolierten Vorhof im Vergleich zur Tyramin- und Nicotinwirkung. Naunyn-Schmiedeberg's Arch. Pharmacol., 242, 458-466.
- 180 Lindmar, R., Löffelholz, K. & Muscholl, E. (1967) Unterschiede zwischen Tyramin und Dimethylphenylpiperazinium in der Ca++-Abhängigkeit und im zeitlichen Verlauf der Noradrenalin-Freisetzung am isolierten Kaninchenherzen. Experientia, 23, 933-934.
- 181 Lindmar, R., Löffelholz, K. & Muscholl, E. (1968) A muscarinic mechanism inhibiting the release of noradrenaline from peripheral adrenergic nerve fibres by nicotinic agents. J. Pharmacol., 32, 280-294.
- 182 Lindmar, R. & Muscholl, E. (1961) Die Wirkung von Cocain, Guanethidin, Reserpin, Hexamethonium, Tetracain und Psicain auf die Noradrenalin-Freisetzung aus dem Herzen. Schmiedeberg's Arch. Pharmacol., 242, 214-227.
- 183 Löffelholz, K. (1967) Untersuchungen über die Noradrenalin-Freisetzung durch Acetylcholin am perfundierten Kaninchenherzen. Naunyn-Schmiedeberg's Arch. Pharmacol., 258, 108-122.
- 184 Löffelholz, K. (1970a) Autoinhibition of nicotinic release of noradrenaline from postganglionic sympathetic nerves. Naunyn-Schmiedberg's Arch. Pharmacol., 267, 49-63
- 185 Löffelholz, K. (1970b) Nicotinic drugs and postganglionic sympathetic transmission. Naunyn-Schmiedeberg's Arch. Pharmacol., 267, 64-73.
- 186 Löffelholz, K. (1975) Muscarinic inhibition of adrenergic neurotransmission. Gordon Conference on Catecholamines, Andover, New Hampshire.
- 187 Löffelholz, K., Lindmar, R. & Muscholl, E. (1967) Der Einfluss von Atropin auf die Noradrenalinfreisetzung durch Acetylcholin. Naunyn-Schmiedeberg's Arch. Pharmacol., 257,308.



- 188 Löffelholz, K. & Muscholl, E. (1969) A muscarinic inhibition of the noradrenaline release evoked by postganglionic sympathetic nerve stimulation. Naunyn-Schmiedeberg's Arch. Pharmacol., 265, 1-15.
- 189 Löffelholz, K. & Muscholl, E. (1970) Inhibition by parasympathetic nerve stimulation of the release of the adrenergic transmitter. Naunyn-Schmiedebergs Arch. Pharmacol., 276, 181-184.
- 190 Loiacono, R.E., Rand, M.J. & Story, D.F. (1985) Interaction between the inhibitory action of acetylcholine and the a-adrenoceptor autoinhibitory feedback system on the release of ³H-noradrenaline from rat atria and rabbit ear artery. Br. J. Pharmacol., 84, 695-705.
- 191 Loiacono, R.E., Rand, M.J. & Story, D.F. (1982) Influence of calcium ion concentration on the interaction of acetylcholine with transmitter noradrenaline release. Clin. Exp. Pharmacol. Physiol., 9, 112-113.
- 192 Loiacono, R.E. & Story D.F. (1984) Acetylchcline induced inhibition of responses to field stimulation in rabbit pulmonary artery is unaffected by endothelium removal. J. Pharm. Pharmacol., 26, 262-264.
- 193 Loiacono, R.E. & Story, D.F. (1986) Effect of α-adrenoceptor agonists and antagonists on cholinergic transmission in guinea-pig isolated atria. Naunyn-Schmiedeberg's Arch.Pharmacol., 334, 40-47.
- 194 Lokhandwala, M.F., Cavero, I.and Buckley, J.P. (1973) Influence of pentobarbital anesthesia on the effects of certain autonomic blocking agents on heart rate. Eur. J. Pharmacol., 24, 274-277.
- 195 Macmillan, W.H. & Rand, M.J. (1962) The effects in rabbits of thyroidectomy and treatment with triiodothyronine on the sensitivity to noradrenaline and the content of noradrenaline in acrta and spleen. J. Pharm. Pharmacol., 14, 257-267.
- 196 Malik, K.U. & Ling, G.M. (1969a) The effect of 1,1-dimethyl-4-phenylpiperazinium on the response of mesenteric arteries to sympathetic nerve stimulation. J. Pharm. Pharmacol. 21, 514-519.
- 197 Malik, K.U. & Ling, G.M. (1969b) Modification by acetylcholine of the response of rat mesenteric arteries to sympathetic stimulation. Circ. Res., 25, 1-9
- 198 Malik, K.U. (1970) Potentiation by antichclinesterases of the response of rat mesenteric arteries to sympathetic postganglionic nerve stimulation. Circ. Res., 27, 647-655
 199 Mathé, A.A., Tong, E.Y and Tisher, P.N. (1977) Norepinephrine
- 199 Mathé, A.A., Tong, E.Y and Tisher, P.N. (1977) Norepinephrine release from the lung by sympathetic nerve stimulation inhibition by the vagus and methacholine. Life Sciences, 20, 1425-1430.
- 200 McCubbin, J.W., Kaneko, Y. & Page, I.H. (1961) The peripheral cardiovascular actions of guanethidine in dogs. J. Pharmacol. Exp. Ther., 131, 346-354.
- 201 McGiff, J.C., Burns, R.B.P. & Blumenthal, M.R. (1967) Role of acetylcholine in the renal vasoconstrictor response to sympathetic nerve stimulation in the dog. Circ. Res., 20, 616-629.
- 202 McNamara, B., Krop, S. & McKay, E.A. (1948) The effect of calcium on the cardiovascular stimulation produced by acetylcholine. J. Pharmacol. Exp. Ther., 92, 153-161.
- 203 Middleton, S., Oberti , C., Prager, R. & Middleton, H. (1956) Stimulating effect of acetylcholine on the papillary myocardium. Acta Physiol. Latinoam., 6, 82-89.

- 204 Millson, D.R. (1959) Nicotine and the effect of antisympathomimetic agents on the aorta of the rabbit. Br. J. Pharmacol., 14, 239-242.
- 205 Mirkin, B.L. & Cervoni, P. (1962) The adrenergic nature of neurohumoral transmission in the cat nictitating membrane following treatment with reserpine. J. Pharmacol. Exp. Ther., 138, 301-308.
- 206 Mottram, D.R., Ivens, C., Lever, J.D. & Presley, R. (1973) A fine structural and enzyme histochemical study of muscle innervation in the normal and 6-hydroxydopamine treated rat vas deferens including a numerical assessment of axonal changes within 48 h of a single injection of the drug. Neuropharmacology, 12, 583-595.
- 207 Muscholl, E. (1970) Cholinomimetic drugs and the release of the adrenergic transmitter. In New Aspects of Storage and Release Mechanisms of Catecholamines, (Ed. H.J.Schümann and G.Kroneberg), pp. 168-186, Springer, Berlin Heidelberg New York.
- 208 Muscholl, E. (1973a) Muscarinic inhibition of the norepinephrine release from peripheral sympathetic fibres. In Pharmacology and the Future of Man. Proceedings of the 5th International Congress of Pharmacology, Vol. 4, Karger, Basel, pp 440-457.
- 209 Muscholl, E. (1973b) Regulation of catecholamine release. The muscarinic inhibitory mechanism. In Frontiers in Catecholamine Research, (Ed. E. Usdin and S. H. Snyder), pp. 537-542, Pergamon Press, New York.
- 210 Muscholl, E. (1979) Presynaptic muscarine receptors and inhibition of release. In The Release of Catecholamines from Adrenergic Neurones, (Ed. D.M.Paton), pp. 87-110, Pergamon, Oxford, New York.
- 211 Muscholl, E. (1980a) Peripheral muscarinic control of norepinephrine release in the cardiovascular system. Am. J. Physiol., 239, H713-H720.
- 212 Muscholl, E. (1980b) From the cholinergic link to the cholinergic antilink in adrenergic transmission: The muscarinic inhibitory mechanism. Trends in Pharmacological Sciences, 1, 381-382.
- 213 Muscholl, E. & Maître, L. (1963) Release by sympathetic stimulation of α -methylnoradrenaline stored in the heart after administration of α -methyldopa. Experientia, 19, 658-659.
- 214 Muscholl, E. & Muth, A. (1982) The effect of physostigmine on the vagally induced muscarinic inhibition of noradrenaline release from the isolated perfused rabbit atria. Naunyn-Schmiedeberg's Arch. Pharmacol., 320, 160-169.
- 215 Muscholl, E., Ritzel, H. & Rossler, K. (1979) Presynaptic muscarinic control of neuronal noradrenaline release. In Presynaptic Receptors, (Ed S.Z.Langer, K.Starke and M.L.Dubocovich), pp. 287-291, Pergamon, Oxford New York.
- 216 Nedergaard, O.A. (1980) Modulation by the muscarinic agonist McN-A-343 of noradrenaline release from vascular sympathetic neurones. J. Cardiovasc. Pharmacol., 2, 629-
- 217 Nedergaard, O.A. & Bevan, J.A. (1969) Effects of nicotine, dimethylphenylpiperazinium and cholinergic blocking agents at adrenergic nerve endings of the rabbit pulmonary artery. J. Pharmacol. Exp. Ther., 168, 127-136.
- 218 Nedergaard, O.A. & Schrold, J.A. (1973) Release of ³H-noradrenaline from incubated and superfused rabbit pulmonary artery. Acta Physiol. Scand., 89, 296.



- 219 Nedergaard, O.A. & Schrold, J.A. (1977) The mechanism of action of nicotine on vascular adrenergic neuroeffector transmission. Eur. J. Pharmacol., 42, 315-329.
- 220 Ng, K.K.F. (1966) The effect of some anticholinesterases on the response of the taenia to sympathetic nerve stimulation. J. Physiol., 182, 233-243.
- 221 Nielsen, K.C., Owman, Ch. & Sporrong, B. (1971) Ultrastructure of the autonomic innervation apparatus in the main pial arteries of rats and cats. Brain Res., 27, 25-32.
- 222 Nystrom, R.A. (1962) Nervous control of the cat nictitating membrane. Am. J. Physiol., 202, 849-854.
- 223 Ohlin, P. & Strömblad, B.C.R. (1963) Observations on the isolated vas deferens. Br. J. Pharmacol., 20, 299-306.
- 224 Pappano, A.J. (1977) Ontogenetic development of autonomic neuroeffector transmission and transmitter reactivity in embryonic and fetal hearts. Pharmacol. Rev., 29, 3-33.
- 225 Patterson, P.H. (1978) Experimental determination of autonomic neurotransmitter functions. Ann. Rev. Neurosci., 1, 1-17.
- 226 Potter, E.K. (1985) Prolonged noradrenergic inhibition of cardiac vagal action following sympathetic stimulation; neuromodulation by neuropeptide Y? Neurosci. Lett., 54, 117-121.
- 227 Potter, E.K. (1987) Cardiac vagal action and plasma levels of neuropeptide Y following intravenous injection in the dog. Neurosci.Lett., 77, 243-247.
- 228 Rand, M.J., McCulloch, M.W. & Story, D.F. (1975) Prejunctional modulation of noradrenergic transmission by noradrenaline, dopamine and acetylcholine. In Central Actions of Drugs in Blood Pressure Regulation, (Ed D.S.Davies and J.L.Reid), pp. 94-132, Pitman Medical, London.
- 229 Rand, M.J. & Stafford, A. (1967) Cardiovascular effects of choline esters. Physiol. Pharmacol., 3, 1-95.
- 230 Rand, M.J. & Varma, B. (1970) The effects of cholinomimetic drugs on responses to sympathetic nerve stimulation and noradrenaline in the rabbit ear artery. Br. J. Pharmacol., 38, 758-770.
- 231 Rand, M.J. & Varma, B. (1971) Effects of the muscarinic agonist McN-A-343 on responses to sympathetic nerve stimulation in the rabbit ear artery. Br. J. Pharmacol., 43, 536-542.
- 232 Rand, M.J. & Wilson, J. (1967) Receptor site of adrenergic neuron blocking drugs. Circ. Res., 20/21 (Suppl. 3), III-89-III-99.
- 233 Richardson, K.C. (1964) The fine structure of the albino rabbit iris with special reference to the identification of adrenergic and cholinergic nerves and nerve endings in its intrinsic muscles. Am. J. Anat., 114, 173-205.
- 234 Richardson, J.A. & Woods, E.F. (1959) Release of norepinephrine from the isolated heart. Proc. Soc. Exp. Biol., N.Y., 100, 149-151.
- 235 Resemblueth, A. & Simeone, F.A. (1934) The interrelationships of vagal and accelerator effects on the cardiac rate. Am. J. Physiol., 110, 42-55.
- Physiol., 110, 42-55.
 236 Ross, G. (1973) Effects of intracoronary infusions of acetylcholine and nicotine on the dog heart in vivo.
 Br.J.Pharmacol., 48, 612-619.
- 237 Samaan, A. (1935) The antagonistic cardiac nerves and heart rate. J. Physiol., 83, 332-340.



- 238 Sarantos-Laska, C., Majewski, H., McCulloch, M.W. & Rand, M.J. (1980a) Mechanism of noradrenaline release from rabbit atria induced by nicotinic agents. Arch. int. Pharmacodyn., 247, 294-305.
- 239 Sarantos-Laska, C., McCulloch, M.W. & Rand, M.J. (1980b)
 Facilitatory effect of nicotine on adrenergic neuroeffector
 transmission in the isolated ear artery of the rabbit. J.
 Pharm. Pharmacol., 32, 92-96.
- Pharm. Pharmacol., 32, 92-96.

 240 Sarantos-Laska, C., McCulloch, M.W., Rand, M.J. & Story, D.F. (1981) Nicotinic release of noradrenaline in the presence of tetrodotoxin from sympathetic nerve terminals in rabbit isolated atria. J. Pharm. Pharmacol., 33, 315-316.
- 241 Sharma, V.K. & Banerjee, S.P. (1977) Presynaptic muscarinic cholinergic receptors in rat heart sympathetic nerves. Eur. J. Pharmacol., 46, 75-76.
- 242 Sharma, V.K. & Banerjee, S.P. (1978) Presynaptic muscarinic cholinergic receptors. Nature, 272, 276-278.
- 243 Sherif, M.A.F. (1935) The chemical transmission of the sympathetic nerve to the uterus. J. Physiol., 85, 298-308.
- 244 Sjöstrand, N.O. (1961) Effect of some smooth muscle stimulants on the motor response of the isolated guinea-pig vas deferens to hypogastric nerve stimulation. Nature, 192, 1190-1191
- to hypogastric nerve stimulation. Nature, 192, 1190-1191 245 Sjöstrand, N.O. & Swedin, G. (1968) Potentiation by smooth muscle stimulants of the hypogastric nerve-vas deferens preparation from normal and castrated guinea-pigs. Acta Physiol. Scand., 74, 472-479.
- 246 Sorimachi, M., Oesch, F. & Thoenen, H. (1973) Effects of colchicine and cytochalasin B on the release of ³H-norepinephrine from guinea-pig atria evoked by high potassium, nicotine and tyramine. Naunyn-Schmiedeberg's Arch. Pharmacol., 276, 1-12.
- 247 Spadolini, I. & Domini, G. (1940) La duplice azione dell'acetilcolina sul cuore isolato di cavia. Arch. Fisiol., 40, 147-172.
- 248 Starke, K. (1977) Regulation of noradrenaline release by presynaptic receptor systems. Rev. Physicl. Biochem. Pharmacol., 77, 1-124.
- 249 Starke, K., Endo, T. & Taube, H.D. (1975c) Relative pre- and post-synaptic potencies of α-adrenoceptor agonists in the rabbit pulmonary artery. Naunyn-Schmiedeberg's Arch. Pharmacol., 291, 55-78.
- 250 Starke, K. & Montel, H. (1974) Influence of drugs with affinity for α -adrenoceptors on noradrenaline release by potassium, tyramine and dimethylphenylpiperazinium. Eur. J. Pharmacol., 27, 273-280.
- 251 Starke, K. & Weitzell, R. (1978) Is histamine involved in the sympathomimetic effect of nicotine? Naunyn-Schmiedeberg's Arch. Pharmacol., 304, 237-248.
- 252 Steinsland, O.S. & Furchgott, R.F. (1975a) Vasoconstriction of the rabbit isolated ear artery caused by nicotinic agonists acting on adrenergic neurones. J. Pharmacol. Exp. Ther., 193, 128-137.
- 253 Steinsland, O.S. & Furchgott, R.F. (1975b) Desensitization of the adrenergic neurones of the isolated rabbit ear artery to nicotinic agonists. J. Pharmacol. Exp. Ther., 193, 138-148.
- 254 Steinsland, O.S., Furchgott, R.F. & Kirpekar, S.M. (1973)Inhibition of adrenergic neurotransmission by parasympathomimetics in the rabbit ear artery. J. Pharmacol. Exp. Ther., 184, 346-356.



- 255 Stjärne, L. (1975) Pre- and post-junctional receptor-mediated cholinergic interactions with adrenergic transmission in the guinea-pig vas deferens. Naunyn-Schmiedeberg's Arch. Pharmacol., 288, 305-310.
- 256 Story, D.F., Allen, G.S., Hope, W., McCulloch, M.W., Rand, M.J. & Sarantos, C (1975) Modulation of adrenergic transmission by acetylcholine. Clin. Exp. Pharmacol. Physiol., 2 (Suppl.) 27-
- 257 Story, D.F., Briley, M.S. & Langer, S.Z. (1979) The effects of 6-hydroxydopamine treatment on the binding of 3H-quinuclidinylbenzilate, 3H-dihydroergocryptine and 3H-WB4101 to rat ventricle membranes. In Presynaptic Receptors (Eds S Z Langer and M L Dubocovich) pp 105-109, Pergamon Press, Oxford, .
- 258 Stuesse, S.L., Wallick, W. & Levy, M.N. (1979) Autonomic control of right atrial contractile strength in the dog. Am. J. Physiol., 236, H860-H865.
- 259 Su, C. & Bevan, J.A. (1970) Blockade of the nicotine-induced norepinephrine release by cocaine, phenoxybenzamine and
- desipramine. J. Pharmacol. Exp. Ther., 175, 533-540. 260 Szentiványi, M. & Kövér, A. (1956) Beeinflussung der zweiphasigen Wirking von Adrenalin und Azetylcholin mit gangliotropen Verbindungen. Acta Physiol. Acad. Sci. Hung., 9, 203-213.
- 261 Taube, H.D., Endo, T., Bangerter, A. & Starke, K. (1976) Presynaptic receptor systems on the noradrenergic nerves of the rabbit pulmonary artery. Naunyn-Schmiedeberg's Arch. Pharmacol., 293, R2.
- 262 Thoenen, H. & Tranzer, J.P. (1968) Chemical sympathectomy by selective destruction of adrenergic nerve endings with 6hydroxydopamine. Naunyn-Schmiedeberg's Arch. Pharmacol., 261, 271-288.
- 263 Thoenen, H., Tranzer, J.P., Hürlimann, S.A. & Haefely, W. (1966) Untersuchungen zur Frage eines cholinergischen Gliedes in der postganglionären sympathischen Transmission. Helv. Physiol. Acta, 24, 229-246.
- 264 Thompson, J.W. (1958) Studies on the responses of the isolated nictitating membrane of the cat. J. Physiol., 141, 46-72.
- 265 Toda, N. (1975) Nicotine-induced relaxation in isolated canine cerebral arteries. J. Pharmacol. Exp. Ther., 193, 376-384.
- 266 Torchiana, M.L. & Angelakos, E.T. (1963) Positive chronotropic and inotropic response of isolated atria to acetylcholine and tyramine. Arch. Intern. Physiol. Biochem. 71, 762-767.
- 267 Uvnas, B. (1960) Sympathetic vasodilator system and blood flow. Physiol. Rev., 40, 69-80.
- 268 Uvnas, B. (1967) Cholinergic vasodilator innervation to skeletal muscles. Circ. Res., 20-21 (Suppl. I) 183.-190.
 269 Van Hee, R. & Vanhoutte, P.M. (1978) Cholinergic inhibition of adrenergic neurotransmission in the canine gastric artery. Gastroenterology, 74, 1266-1270.
- 270 Vanhoutte, P.M. (1974) Inhibition by acetylcholine of adrenergic neurotransmission in vascular smooth muscle. Res., 34, 317-326.
- 271 Vanhoutte, P.M. (1977) Cholinergic inhibition of adrenergic transmission. Fed. Proc., 36, 2444-2449.
- 272 Vanhoutte, P.M. & Levy, M.N. (1980) Prejunctional modulation of adrenergic neurotransmission in the cardiovascular system. Am. J. Physiol., 238, H275-H281.



- 273 Vanhoutte, F.M., Lorenz, R.R. & Tyce, G.M. (1973) Inhibition of norepinephrine-3H release from sympathetic nerve endings in veins by acetylcholine. J. Pharmacol. Exp. Ther., 185, 386-394.
- 274 Vanhoutte, P.M. & Shepherd, J.T. (1973) Venous relaxation caused by acetylcholine acting on the sympathetic nerves. Circ. Res., 32, 259-267.
- Vanhoutte, P.M. & Verbeuren, T.J. (1975) Inhibition of norepinephrine release in vascular smooth muscle by acetylcholine. Due to hyperpolarization of adrenergic nerve endings? Arch. int. Pharmacodyn., 213, 332-333.
 Vanhoutte, P.M. & Verbeuren, T.J. (1976a) Inhibition by
- 276 Vanhoutte, P.M. & Verbeuren, T.J. (1976a) Inhibition by acetylcholine of the norepinephrine release evoked by potassium in canine saphenous veins. Circ. Res., 39, 263-269
- 277 Vanhoutte, P.M. & Verbeuren, T.J. (1976b) Inhibition by acetylcholine of ³H-norepinephrine release in cutaneous veins after alpha-adrenergic blockade. Arch. int. Pharmacodyn. 221, 344-346
- 278 Vanhoutte, P.M., Verbeuren, T.J. & Collis, M.G. (1977)
 Muscarinic inhibition of adrenergic neurotransmission is not due to a decrease in Ca⁺⁺ entry into the adrenergic nerve endings. J. Pharmacol. (Paris), 8, 556-557.
- 279 Vanhoutte, P.M. Verbeuren, T.J. & Webb, R.C. (1982) Local modulation of adrenergic neuroeffector interaction in the blood vessel wall. Physiol. Rev., 61, 151-247.
- 280 Varagic, V. (1956) An isolated rabbit hypogastric-nerve-uterus preparation, with observations on the hypogastric transmitter. J. Physiol., 132, 92-99.
- 281 Verbeuren, T.J. & Vanhoutte, P.M. (1976) Acetylcholine inhibits potassium evoked release of ³H-norepinephrine in different blood vessels of the dog. Arch. int. Pharmacodyn., 221. 347-350.
- 221, 347-350. 282 Vizi, E.S. (1979) Presynaptic modulation of neurochemical transmission. Prog. Neurobiol., 12, 181-290.
- 283 Vizi, E.S. & Knoll, J. (1971) The effects of sympathetic nerve stimulation and guanethidine on parasympathetic neuroeffector transmission; the inhibition of acetylcholine release. J. Pharm. Pharmacol., 23, 918-925.
- 284 von Euler, U.S. & Gaddum, J.H. (1931) Pseudomotor contractures after degeneration of the facial nerve. J. Physiol., 73, 54-66.
- 285 Wallick, D.W., Felder, D. & Levy, M.N. (1978) Autonomic control of pacemaker activity in the atrioventricular junction of the dog. Am. J. Physiol., 235, H308-H313.
- 286 Wallick, D.W., Martin, P.J., Masuda, Y. & Levy, M.N. (1982)
 Effects of autonomic activity and changes in heart rate on
 atrioventricular conduction. Am.J. Physiol., 243, H523-H527.
- atrioventricular conduction. Am.J. Physiol., 243, H523-H527. 287 Warner, H.R. & Russell, R.O. (1969) Effect of combined sympathetic and vagal stimulation on heart rate in the dog. Circ. Res., 24, 567-573.
- 288 Watanabe, A.M., Lindemann, J.P. & Fleming, J.W. (1984)
 Mechanisms of muscarinic modulation of protein phosphorylation
 in intact ventricles. Fed. Proc., 43, 2618-2623.
- 289 Waterson, J.G., Hume, W.R. & de la Lande, I.S. (1970) The distribution of cholinesterase in the rabbit ear artery. J. Histochem. Cytochem., 18, 211.
- 290 Weiner, N. (1980) The role of cyclic nucleotides in the regulation of neurotransmitter release from adrenergic neurons by neuromodulators. In Neurochemistry and Neuropharmacology, Vol. 4, (Ed Youdin, M.B.H., Lovenberg, W., Sharman, D.F. & Lagnado, J.R.), pp. 69-124. John Wiley & Sons.



- 292 Westfall, T.C. (1971b) Studies on the mechanism of action of nicotinic agents on adrenergic nerve terminals.

 Pharmacologist. 13, 229
- 293 Westfall, T.C. (1977) Local regulation of adrenergic neurotransmission. Physiol. Rev., 57, 659-728.
- 294 Westfall, T.C. & Brasted, M. (1972) The mechanism of action of nicotine on the adrenergic neurones in the perfused guinea-pig heart. J. Pharmacol. Exp. Ther., 182, 409-418.
- 295 Westfall, T.C. & Brasted, M. (1973) Effect of 4,4'-biphenylenebis-[(2-oxoethylene)-bis-(2,2-diethoxyethyl)]dimethylammonium dibromide (DMAE) on accumulation and nicotine-induced release of norepinephrine in the heart. J. Pharmacol. Exp. Ther., 184, 198-204.
- 296 Westfall, T.C. & Brasted, M. (1974) Specificity of blockade of the nicotine-induced release of ³H-norepinephrine from adrenergic neurons of the guinea-pig heart by various pharmacological agents. J. Pharmacol. Exp. Ther., 189, 659-664.
- 297 Westfall, T.C. & Hunter, P.E. (1974) Effect of muscarinic agonists on the release of ³H-noradrenaline from the guineapig perfused heart. J. Pharm. Pharmacol., 26, 458-460.
- 298 Westfall, T.C. & Saunders, J. (1977) Absence of nicotinic receptors mediating the release of norepinephrine from adrenergic neurons in the rat heart. Proc. Soc. Exp. Biol. Med., 156, 476-479.
- 299 Wilson, A.B. (1962) Adrenergic neurone blocking action of dimethylphenylpiperazinium. J. Pharm. Pharmacol., 14, 700.
- 300 Wetzel, G.T. Goldstein, D. & Brown, J.H. (1985) Acetylcholine release from rat atria can be regulated through an α_1 -adrenergic receptor. Circ. Res., 56, 763-766.

